

Updates of W_R effects on CP angles determination in B decays

H. Hayakawa¹, K. Hosokawa² and T. Kurimoto³,
 Department of Physics, Faculty of Science,
 Toyama University,
 Toyama 930, Japan

Abstract

The recently observed CP violation in B decay and B - $\bar{B}$ mixing data put constraints on the mass of W_R and the parameters of the right-handed current quark mixing matrix V^R in $SU(2)_L \times SU(2)_R \times U(1)$ gauge model. It is shown that the allowed region of parameters are severely restricted for light W_R with mass on the order of 1 TeV. There exist sets of parameters which can accommodate large CP violation as measured by Belle, $\sin 2\phi_1|_{exp} \simeq 1$, for $M_{W_R} = 1 \sim 10$ TeV.

¹e-mail: haya@jodo.sci.toyama-u.ac.jp

²e-mail: kaoru@jodo.sci.toyama-u.ac.jp

³e-mail: krmt@k2.sci.toyama-u.ac.jp

The B factories at KEK and SLAC have established the existence of CP violation in B meson system by measuring the time-dependent CP asymmetry[1] of neutral B meson decays into $(c\bar{c})$ meson+neutral $K^{(*)}$ meson ;

$$A(t) = \frac{\Gamma[\overline{B}^0(t) \rightarrow f_{CP}] - \Gamma[B^0(t) \rightarrow f_{CP}]}{\Gamma[\overline{B}^0(t) \rightarrow f_{CP}] + \Gamma[B^0(t) \rightarrow f_{CP}]} = -\xi_f \sin 2\phi_1 \sin(\Delta M_B t), \quad (1)$$

where ξ_f is the CP eigenvalue of the final state. They obtained[2, 3]

$$\sin 2\phi_1 = \begin{cases} 0.59 \pm 0.14 \pm 0.05 & (BABAR) \\ 0.99 \pm 0.14 \pm 0.06 & (Belle) \end{cases}. \quad (2)$$

Let us check if the above value is consistent with the 3-generation standard model with Kobayashi-Maskawa mechanism of CP violation[4]. With the notation of the unitarity




Figure 1: Unitarity triangle

triangle given in Fig.1 where $V_{cb}^*V_{cd}$ is real negative, geometrically defined $\sin 2\phi_1$ is given as

$$\sin 2\phi_1 = \sin \left(2\pi - 2 \arg \left[\frac{-V_{tb}^*V_{td}}{-V_{cb}^*V_{cd}} \right] \right) = \sin \left(2 \arg \left[-1 + \left| \frac{V_{ub}^*V_{ud}}{V_{cb}^*V_{cd}} \right| e^{-i\phi_3} \right] \right). \quad (3)$$

If no new physics beyond the 3-generation standard model enters in the measured processes of CP violation, the observed $\sin 2\phi_1$ should agree with the above geometrically defined one. Assuming that new physics does not affect the determination of $|V_{ud}|$, $|V_{cd}|$ and $|V_{ub}/V_{cb}|$ which are obtained through tree level semi-leptonic processes, the prediction of $\sin 2\phi_1$ in terms of ϕ_3 by using eq.(3) and $|(V_{ub}^*V_{ud})/(V_{cb}^*V_{cd})|$ is given as Fig.2. The measured result $\sin 2\phi_1 > 0.4$ is consistent with $\phi_3 = 15^\circ \sim 145^\circ$. The neutral B meson mass difference ΔM_B in the standard model is also estimated as a function of ϕ_3 . Once $|V_{ub}/V_{cb}|$ is given, $V_{tb}V_{td}^*$ can be expressed in terms of ϕ_3 and $|V_{ub}/V_{cb}|$ by using unitarity. Taking the ambiguity of hadron matrix elements to $\pm 30\%$ and errors of $|V_{ub}/V_{cb}|$, we find the standard model is consistent with the experimental value of ΔM_B for $\phi_3 = 20^\circ \sim 70^\circ$.

Thus, the measured values of $\sin 2\phi_1$ are consistent with the 3-generation standard model considering the errors at present, though the central value given by Belle cannot be attained in the standard model. If the high value by Belle is confirmed in the future experiments, some new physics beyond the standard model is necessary.

In this work we investigate $SU(2)_L \times SU(2)_R \times U(1)$ model (L-R model)[5] as a possible candidate of new physics which can give larger CP asymmetry in the $\sin 2\phi_1$ determination than in the standard model. We investigate constraints on the model,




Figure 2: $\sin 2\phi_1$ in the 3-generation standard model. Upper, middle and lower curves correspond to $|V_{ub}/V_{cb}| = 0.11$, 0.09 and 0.07 , respectively.




Figure 3: $\Delta M_B^{SM}/\Delta M_B^{exp}$ in the 3-generation standard model. Upper, middle and lower curves correspond to $|V_{ub}/V_{cb}| = 0.07$, 0.09 and 0.11 , respectively.

and explore the possibility of obtaining a high value of CP asymmetry simultaneously satisfying constraints by ΔM_B and $K-\bar{K}$ system. Some groups including one of the present authors have investigated L-R model and showed that the gauge boson coupled to right-handed charged current (W_R) can affect significantly on the determination of the CP violation angles in B decays[6, 7, 8, 9]. The essence is as follows; Though W_R is much heavier than the ordinary W boson, some elements of the right-handed current quark mixing matrix V^R are not necessary suppressed in comparison with the CKM mixing matrix elements[4, 10]. Then W_R can contribute significantly to some processes like $B-\bar{B}$ mixing where ordinary W boson contribution is much CKM suppressed.

There exists a sizable contribution to $K-\bar{K}$ mixing in L-R model from the box diagram with one W and one W_R exchange[6, 11, 12, 13], which allows only following forms of V^R to avoid constraint from CP violation in $K-\bar{K}$ mixing for not too heavy W_R with mass of $O(1)$ TeV or less [8];

$$V_I^R = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & e^{i\omega} \end{pmatrix}, \quad V_{II}^R = \begin{pmatrix} 0 & 1 & 0 \\ \cos \theta_R & 0 & -e^{i\omega} \sin \theta_R \\ \sin \theta_R & 0 & e^{i\omega} \cos \theta_R \end{pmatrix}. \quad (4)$$

(If we allow fine tunings among the parameters of CKM matrix and those of V^R , there are other possibilities, which we do not consider here.) The former, V_I^R , does not give significant contribution to either $B-\bar{B}$ mixing or b decay, so we concentrate on the latter type of V^R here.

The contribution to $B-\bar{B}$ mixing is written as

$$M_{12}^B = M_{12}^{SM} + M_{LR} + M_{RR}, \quad (5)$$

where M_{12}^{SM} is the standard model contribution, M_{LR} from the box diagram with one W and one W_R exchange, and M_{RR} from two W_R exchange. M_{RR} is obtained simply by exchanging $L \leftrightarrow R$ in the standard model contribution. M_{LR} is calculated from the following effective Hamiltonian[7, 15, 16];

$$\mathcal{H}_{LR}^{eff} = \sum_{i,j=u}^t \frac{2G_F^2 M_W^2}{\pi^2} \beta_g V_{id}^{L*} V_{ib}^R V_{jd}^{R*} V_{jb}^L J(x_i, x_j, \beta) \bar{d}_R b_L \bar{d}_L b_R + (\text{h.c.}), \quad (6)$$

where $\beta_g = (g_R/g_L)^2 (M_W^2/M_{W_R}^2)$ and $x_i = m_i^2/M_W^2$. The loop function is defined as

$$J(x, y, \beta) \equiv \sqrt{xy} [(\eta^{(1)} + \eta^{(2)} \frac{xy\beta}{4}) J_1(x, y, \beta) - \frac{1}{4}(\eta^{(3)} + \eta^{(4)}\beta) J_2(x, y, \beta)], \quad (7)$$

with

$$J_1(x, y, \beta) = \frac{x \ln x}{(1-x)(1-x\beta)(x-y)} + (x \leftrightarrow y) - \frac{\beta \ln \beta}{(1-\beta)(1-x\beta)(1-y\beta)},$$

$$J_2(x, y, \beta) = \frac{x^2 \ln x}{(1-x)(1-x\beta)(x-y)} + (x \leftrightarrow y) - \frac{\ln \beta}{(1-\beta)(1-x\beta)(1-y\beta)},$$

where $\eta^{(1)-(4)}$ are QCD corrections. We use here $\eta^{(1)} = 1.1$, $\eta^{(2)} = 0.26$, $\eta^{(3)} = 1.1$, $\eta^{(4)} = 1.0$ as the values of QCD corrections[14].




Figure 4: Allowed region by ΔM_B (left) and $Asy(\Psi K)$ (right) for $M_{W_R} = 1$ TeV and $\phi_3 = 135^\circ$. Black painted regions are consistent with experimental value of ΔM_B in the left figure.




Figure 5: $\Delta M_B|_{theory}/\Delta M_B|_{exp}$ and $Asy(\Psi K)$ for $M_{W_R} = 1$ TeV, $\phi_3 = 135^\circ$ and $\theta_R = 100^\circ$.

Now we evaluate M_{12}^B varying θ_R and ω in V^R with the following inputs; $M(W_R) = 1 \sim 10$ TeV, ϕ_3 in $V_{KM} = 45^\circ, 90^\circ, 135^\circ$, $|V_{ub}/V_{cb}| = 0.09$, $f_B\sqrt{B_B} = 230$ MeV. Then we draw allowed regions by the experimental values of ΔM_B allowing $\pm 30\%$ ambiguity from errors in $f_B\sqrt{B_B}$ and $|V_{ub}/V_{cb}|$, and estimate CP asymmetry in $B \rightarrow (c\bar{c}) + K^{(*)}$ corresponding to $\sin 2\phi_1$, which we call as $Asy(\Psi K)$. First we take $M_{W_R} = 1$ TeV and $\phi_3 = 135^\circ$. The allowed region and the predicted $Asy(\Psi K)$ are shown in Fig.4. It can be seen that only small portions of parameter space in θ_R and ω are allowed by ΔM_B . We fix $\theta_R = 100^\circ$ and estimate ΔM_B and $Asy(\Psi K)$. With $\theta_R = 100^\circ$ the CP phase ω in V^R is restricted to $30^\circ \sim 90^\circ$. If we further impose $Asy(\Psi K) > 0.4$ from the recent measurement, ω should be less than 60° . It is interesting that large CP asymmetry given by Belle, $Asy(\Psi K) \sim 1$, is possible for $\omega = 30^\circ \sim 45^\circ$. Similar figures for $\theta_R = 90^\circ$ and 45° are shown in Figs.6 and 7. The allowed region of θ_R and ω is severely restricted. This is because W_R gives significant contribution to B - $\bar{B}$ mixing even for $M_{W_R} \sim 1$ TeV as pointed out in ref.[8]. The standard model contribution M_{12}^{SM} is CKM suppressed by




Figure 6: Same as Fig.4 for $\phi_3 = 90^\circ$.




Figure 7: Same as Fig.4 for $\phi_3 = 45^\circ$.

λ^6 ($\lambda \equiv |V_{us}| = 0.22$) while M_{LR} by λ^3 . Though another suppression of $(M_W/M_{W_R})^2$ is incorporated in M_{LR} , the enhancement in loop function and λ^{-3} factor make M_{LR} similar order of magnitude with M_{12}^{SM} .

We have made same calculations for $M_{W_R} = 2, 3, 5$ and 10 TeV. The results are shown in Figs.8-16 and 18. The area of allowed region becomes maximal at $M_{W_R} = 3$ TeV for $\phi_3 = 135^\circ$. The standard model contribution M_{12}^{SM} alone cannot give ΔM_B consistent with the experimental data for $\phi_3 = 135^\circ$. With suitable magnitude of W_R contribution we can give experimentally consistent value of ΔM_B . Too heavy W_R cannot give sufficient contribution to compensate for M_{12}^{SM} . Similar situation occurs for $\phi_3 = 90^\circ$ at more larger M_{W_R} . No allowed region remains at $M_{W_R} = 10$ TeV for $\phi_3 = 90^\circ$ and 135° . The allowed region spreads as M_{W_R} gets larger for $\phi_3 = 45^\circ$ since the standard model contribution M_{12}^{SM} alone gives ΔM_B and $Asy(\Psi K)$ consistent with experimental data for $\phi_3 = 45^\circ$. It is interesting that not a small are of allowed regions which give large CP asymmetry remains even for heavy W_R . For example, the figure for $M_{W_R} = 5$ TeV, $\phi_3 = 45^\circ$ and $\theta_R = 30^\circ$ is shown in Fig.17, and the figure for $M_{W_R} = 10$ TeV, $\phi_3 = 45^\circ$ is shown in Fig.18.

Let us comment on other CP angles, ϕ_2 and ϕ_3 . ϕ_2 is measured in the CP asymmetry in $B \rightarrow \pi\pi$ decay. CP violation occurs through the interference among B - $\bar{B}$ mixing, tree and penguin decays of $b \rightarrow u\bar{u}d$. W_R can contribute significantly to B - $\bar{B}$ mixing as in the case of ϕ_1 . There also exists contribution to $b \rightarrow d$ penguin. The ratio to the standard




Figure 8: Same as Fig.4 for $M_{W_R} = 2$ TeV and $\phi_3 = 135^\circ$.




Figure 9: Same as Fig.4 for $M_{W_R} = 2$ TeV and $\phi_3 = 90^\circ$.




Figure 10: Same as Fig.4 for $M_{W_R} = 2$ TeV and $\phi_3 = 45^\circ$.




Figure 11: Same as Fig.4 for $M_{W_R} = 3$ TeV and $\phi_3 = 135^\circ$.




Figure 12: Same as Fig.4 for $M_{W_R} = 3$ TeV and $\phi_3 = 90^\circ$.




Figure 13: Same as Fig.4 for $M_{W_R} = 3$ TeV and $\phi_3 = 45^\circ$.




Figure 14: Same as Fig.4 for $M_{W_R} = 5$ TeV and $\phi_3 = 135^\circ$.




Figure 15: Same as Fig.4 for $M_{W_R} = 5$ TeV and $\phi_3 = 90^\circ$.




Figure 16: Same as Fig.4 for $M_{W_R} = 5$ TeV and $\phi_3 = 45^\circ$.




Figure 17: $\Delta M_B|_{theory}/\Delta M_B|_{exp}$ and $Asy(\Psi K)$ for $M_{W_R} = 5$ TeV, $\phi_3 = 45^\circ$ and $\theta_R = 30^\circ$.




Figure 18: Same as Fig.4 for $M_{W_R} = 10$ TeV and $\phi_3 = 45^\circ$.




Figure 19: $\Delta\phi_3$ for $M_{W_R} = 1$ TeV, $\phi_3 = 135^\circ$ and $\theta_R = 100^\circ$.

model penguin up to log loop function is given as

$$\frac{g_L^2}{M_W^2} V_{tb}^* V_{td} : \frac{g_R^2}{M_{W_R}^2} V_{tb}^{R*} V_{td}^R = 1 : \beta_g \frac{e^{-i\omega} \sin 2\theta_R}{2V_{tb}^* V_{td}}. \quad (8)$$

The magnitude of $|\beta_g/(2V_{tb}^* V_{td})|$ is about 0.3 for $M_{W_R} = 1$ TeV. W_R penguin is less than 10% of the standard model one taking the allowed region of θ_R into account. So we can neglect $b \rightarrow d W_R$ penguin. Then the effect on ϕ_2 is same as ϕ_1 . For ϕ_3 we consider the measurement by using $B^\pm \rightarrow DK$. CP violation occurs through the interference between tree decays, $\bar{b} \rightarrow \bar{c} u \bar{s}$ and $\bar{b} \rightarrow \bar{u} c \bar{s}$ with common final state. W_R does not contribute to $\bar{b} \rightarrow \bar{u} c \bar{s}$ as $V_{ub}^R = 0$, but can affect $\bar{b} \rightarrow \bar{c} u \bar{s}$ decay.

$$\frac{g_L^2}{M_L^2} V_{cb}^* V_{us} : \frac{g_R^2}{M_R^2} V_{cb}^{R*} V_{us}^R = 1 : \beta_g \frac{(-e^{-i\omega} \sin \theta_R)}{V_{cb} V_{us}}. \quad (9)$$

The deviation of measured ϕ_3 , $\Delta\phi_3$ from the standard model value for $M_{W_R} = 1$ TeV, $\phi_3 = 135^\circ$ in CKM matrix and $\theta_R = 100^\circ$ is given in Fig.19. The deviation can reach -45° for $\omega = 40^\circ$. As W_R gets heavier the deviation becomes small in proportional to $1/M_{W_R}^2$. This deviation cannot be observed in the measurements of ϕ_3 in $B \rightarrow K\pi$ since $V_{us}^R V_{ub}^{R*} = 0$. So we can expect disagreement between two kinds of measurements of ϕ_3 .

In conclusion, we have investigated W_R effects on B - $\bar{B}$ mixing and CP asymmetry in B decays, and found that W_R effect is sizable even for $M_{W_R} = 1 \sim 10$ TeV. The experimental

values of ΔM_B and CP asymmetry in $B \rightarrow (c\bar{c}) + K^{(*)}$, $Asy(\Psi K)$, severely constrain the parameters of right-handed quark mixing matrix V^R . With allowed parameters the CP asymmetry $Asy(\Psi K)$ can be as large as 1 which is the central value of Belle. If future experiments confirms the high value of $Asy(\Psi K)$, fine measurements of ϕ_2 and ϕ_3 in various modes are necessary to distinguish this kind of model and other new physics.

Acknowledgments

This work is supported in part by Grant-in Aid for Scientific Research from the Ministry of Education, Science and Culture of Japan under the Grant No.11640265.

References

- [1] A.B. Carter and A.I. Sanda, Phys. Rev. Lett. **45** 952 (1980), Phys. Rev. **D23** 1567 (1981);
I.I. Bigi and A.I. Sanda, Nucl. Phys. **B193** 85 (1981), Nucl. Phys. **B281** 41 (1987).
- [2] BABAR Collaboration, Phys. Rev. Lett. **87**, 091802 (2001).
- [3] Belle Collaboration, Phys. Rev. Lett. **87**, 091801 (2001).
- [4] M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
- [5] R.N. Mohapatra and J.C. Pati, Phys. Rev. **D11** 566 (1975), **D11** 2558 (1975);
G. Senjanovic and R.N. Mohapatra, Phys. Rev. **D12** 1502 (1975);
G. Senjanovic, Nucl. Phys. **B153** 334 (1979).
- [6] D. London and D. Wyler, Phys. Lett. **B232** 503 (1989).
- [7] H. Nishiura, E. Takasugi and M. Tanaka, Prog.Theor.Phys. **84**, 116 (1990); **84**, 502 (1990);
85, 343 (1991).
- [8] T. Kurimoto, A. Tomita and S. Wakaizumi, Phys. Lett. **B381**, 470 (1996).
- [9] D. Silverman and H. Yao, hep-ph/9706359 .
- [10] N. Cabibbo, Phys. Rev. Lett, **10**, 531 (1963).
- [11] F.I. Olness and M.E. Ebel, Phys. Rev. **D30** 1034 (1984).
- [12] P. Langacker and S.U. Sankar, Phys. Rev. **D40** 1569 (1989).
- [13] G. Barenboim, J. Bernabeu, J. Prades, M. Raidal, Phys.Rev. **D55**, 4213 (1997).
- [14] A. Datta, E.A. Paschos, J.-M.Schwartz and M.N.Sinha Roy, Z.Phys. **C46**, 63 (1990); hep-ph/9509420 (1995), DO-TH 95/12.
- [15] G. Beall, M. Bander and A. Soni, Phys. Rev. Lett. **48** 848 (1982).
- [16] G. Ecker and W. Grimus, Nucl. Phys. **B258** 328 (1985);